

SUMS OF POWERS OF CONSECUTIVE q -INTEGERS

TAEKYUN KIM

Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr

ABSTRACT. Let n, k be the positive integers ($k > 1$), and let $S_{n,q}(k)$ be the sums of the n -th powers of positive q -integers up to $k-1$: $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l [l]_q^n$, where $[l]_q = \frac{q^l - 1}{q - 1}$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$ as follows:

$$S_{n+1,q}(k) = \frac{1}{n+1} \sum_{i=0}^n \binom{n+1}{i} \beta_{i,q} q^{ki} [k]_q^{n+1-i} + \frac{(1 - q^{(n+1)k}) \beta_{n+1,q}}{n+1},$$

where $\beta_{i,q}$ are the q -Bernoulli numbers.

1. INTRODUCTION

Let n be a natural numbers. There are formulas such as

$$\begin{aligned} 1 + 2 + \cdots + n &= \frac{n^2 + n}{2}, \\ 1^2 + 2^2 + \cdots + n^2 &= \frac{2n^3 + 3n^2 + n}{6}, \\ 1^3 + 2^3 + \cdots + n^3 &= \frac{n^4 + 2n^3 + n^2}{4}, \dots \end{aligned}$$

In Korea, these are subjects of the high school mathematics. J. Bernoulli (1713) first discovered the method which one can produce those formulae for the sum $\sum_{l=1}^n l^k$ for

2000 Mathematics Subject Classification 11S80, 11B68

Key words and phrases: q -Bernoulli number, p -adic q -integrals, zeta function, Dirichlet series

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

any natural numbers. Let q be an indeterminate which can be considered in complex number field, and for any integer k define the q -integer as $[k]_q = \frac{q^k - 1}{q - 1}$, cf. [1,2]. Note that $\lim_{q \rightarrow 1} [k]_q = k$. In this paper we evaluate sums of powers of consecutive q -integers. For any positive integers $n, k (> 1)$, let $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l [l]_q^n$. Following an idea due to J. Bernoulli, we explore a formula for $S_{n,q}(k)$ as follows:

$$S_{n+1,q}(k) = \frac{1}{n+1} \sum_{i=0}^n \binom{n+1}{i} \beta_{i,q} q^{ki} [k]_q^{n+1-i} + \frac{(1 - q^{(n+1)k}) \beta_{n+1,q}}{n+1},$$

where $\beta_{i,q}$ are the q -Bernoulli numbers.

2. SUM OF THE n -TH POWERS OF POSITIVE q -INTEGERS UP TO $k - 1$

Let j be the positive integers. Then we easily see that

$$(1) \quad [j+1]_q^2 - [j]_q^2 = ([j]_q + q^j)^2 - [j]_q^2 = q^j (2[j]_q + q^j).$$

By (1), we have

$$(2) \quad [k]_q^2 = \sum_{j=0}^{k-1} ([j+1]_q^2 - [j]_q^2) = 2 \sum_{j=0}^{k-1} q^j [j]_q + \frac{[2k]_q}{[2]_q}.$$

Hence,

$$(3) \quad \sum_{j=0}^{k-1} q^j [j]_q = \left([k]_q^2 - \frac{[2k]_q}{[2]_q} \right) \frac{1}{2}.$$

By the same method of Eq.(1), we easily see that

$$(4) \quad [j+1]_q^3 - [j]_q^3 = 3[j]_q^2 q^{j+1} + 3[j]_q q^j + q^{3j}.$$

Thus we have

$$(5) \quad \sum_{j=0}^{k-1} [j]_q^2 q^{j+1} = \frac{1}{3} [k]_q^3 - \frac{1}{2} \left([k]_q^2 - \frac{[2k]_q}{[2]_q} \right) - \frac{1}{3} \frac{[3k]_q}{[3]_q}.$$

Example. For $q = \frac{9}{10}$ in (5), we note that

$$\begin{aligned} & \left(\frac{9}{10}\right)^2 + \left(\frac{9}{10}\right)^3 \left(1 + \frac{9}{10}\right)^2 + \cdots + \left(\frac{9}{10}\right)^k \left(1 + \frac{9}{10} + \cdots + \left(\frac{9}{10}\right)^{k-2}\right)^2 + \cdots \\ &= \frac{1}{3} \left(\frac{1}{1 - \frac{9}{10}}\right)^3 - \frac{1}{2} \left(\left(\frac{1}{1 - \frac{9}{10}}\right)^2 - \frac{1}{1 - (\frac{9}{10})^2}\right) - \frac{1}{3} \left(\frac{1}{1 - (\frac{9}{10})^3}\right). \end{aligned}$$

Let n, k be the positive integers ($k > 1$) and let $S_{n,q}(k) = \sum_{l=0}^{k-1} q^l [l]_q^n$. Then we note that

$$(6) \quad \sum_{i=0}^{n-1} \binom{n}{i} S_{i,q^{n-i}}(k) = \sum_{l=0}^{k-1} ([l+1]_q^n - [l]_q^n) = [k]_q^n.$$

By replacing n by $n+1$, we see that

$$(7) \quad [k]_q^{n+1} = \sum_{i=0}^n \binom{n+1}{i} S_{i,q^{n+1-i}}(k) = \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q^{n+1-i}}(k) + (n+1) S_{n,q}(k).$$

Therefore we obtain the following:

$$(8) \quad (n+1) S_{n,q}(k) = [k]_q^{n+1} - \sum_{i=0}^{n-1} \binom{n+1}{i} S_{i,q^{n+1-i}}(k).$$

3. q -ANALOGS OF BERNOULLI POLYNOMIALS

In this section, we assume $q \in \mathbb{C}$ with $|q| < 1$. The q -Bernoulli polynomials $\beta_{n,q}(x)$ are defined by means of the generating function $F_q(t)$ as follows:

$$(9) \quad F_q(t) = e^{\frac{t}{1-q}} \frac{q-1}{\log q} - t \sum_{n=0}^{\infty} q^{n+x} e^{[n+x]_q t} = \sum_{n=0}^{\infty} \frac{\beta_{n,q}(x)}{n!} t^n, \quad |q| < 1, |t| < 1.$$

In the case $x = 0$, $\beta_{n,q}(= \beta_{n,q}(0))$ will be called the q -Bernoulli numbers, cf. [1, 2]. By (9), we easily see that

$$(10) \quad \beta_{n,q}(x) = \sum_{j=0}^n \binom{n}{j} q^{jx} \beta_{j,q} [x]_q^{n-j} = \left(\frac{1}{1-q}\right)^n \sum_{k=0}^n \binom{n}{k} \frac{k}{[k]_q} q^{kx} (-1)^k.$$

In (9), (10), the q -Bernoulli numbers can be rewritten as

$$(11) \quad \beta_{0,q} = \frac{q-1}{\log q}, \quad (q\beta + 1)^k - \beta_{k,q} = \delta_{k,1},$$

where $\delta_{k,1}$ is Kronecker symbol and we use the usual convention about replacing β^i by $\beta_{i,q}$, cf. [1,2]. By (9), (10), and (11), we easily see that

$$-\sum_{l=0}^{\infty} q^{l+k} e^{[l+k]_q t} + \sum_{l=0}^{\infty} q^l e^{[l]_q t} = \sum_{n=1}^{\infty} \left(n \sum_{l=0}^{k-1} q^l [l]^{n-1} \right) \frac{t^{n-1}}{n!}.$$

Thus we have

$$(12) \quad \beta_{n,q}(k) - \beta_{n,q} = n \sum_{l=0}^{k-1} q^l [l]_q^{n-1}.$$

By (12), we obtain the following:

$$(13) \quad \sum_{l=0}^{k-1} q^l [l]_q^n = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} q^{kl} \beta_{l,q} [k]_q^{n+1-l} + \frac{(1 - q^{(n+1)k})}{n+1} \beta_{n+1,q}.$$

In Eq.(10), we know that

$$(14) \quad \beta_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} \beta_{k,q} \sum_{l=0}^k \binom{k}{l} [x]_q^l (q-1)^l.$$

By (14), we easily see that

$$(15) \quad \int_0^k \beta_{n,q}(x) d[x]_q = \frac{1}{n+1} (\beta_{n+1,q}(k) - \beta_{n+1,q}) = S_{n,q}(k).$$

REFERENCES

1. T. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Math., Springer-Verlag, New York, 1986.
2. T. Kim, *q-Volkenborn Integration*, Russian J. Math. Phys. **9** (2002), 288-299.